



Consulting, help, relaxation

INTERNATIONAL JOURNAL OF ENGINEERING SCIENCES & MANAGEMENT

HEALTH SEEKING BEHAVIOUR AND HEALTH SERVICE UTILIZATION IN LUCKNOW

Siraj Ahmad¹ and Adeel Maqbool²

1, Research Scholar, Mewar University, Rajasthan, India

2, Naravadeshwar Management College, Lucknow, UP, India

ABSTRACT

There is a growing literature on health seeking behaviors and the determinants of health services utilization especially in the context of developing countries. However, very few focused studies have been seen in Lucknow in this regard. This paper presents an extensive literature review of the situation in developing countries and relates the similar factors responsible for shaping up of a health seeking behavior and health service utilization in Lucknow. The factors determining the health behaviors may be seen in various contexts: physical, socio-economic, cultural and political. Therefore, the utilization of a health care system, public or private, formal or non-formal, may depend on socio-demographic factors, social structures, level of education, cultural beliefs and practices, gender discrimination, status of women, economic and political systems environmental conditions, and the disease pattern and health care system itself. Policy makers need to understand the drivers of health seeking behavior of the population in an increasingly pluralistic health care system. Also a more concerted effort is required for designing behavioral health promotion campaigns through inter-sector collaboration focusing more on disadvantaged segments of the population.

Key words: Health, health care system, health seeking behavior, health service utilization

INTRODUCTION

Strategic policy formation in all health care systems should be based on information relating to health promoting, seeking and utilization behavior and the factors determining these behaviors. All such behaviors occur within some institutional structure such as family, community or the health care services. The factors determining the health behaviors may be seen in various contexts: physical, socio-economic, cultural and political. Therefore, the utilization of a health care system, public or private, formal or non-formal, may depend on socio-demographic factors, social structures, level of education, cultural beliefs and practices, gender discrimination, status of women, economic and political systems environmental conditions, and the disease pattern and health care system itself. A main driver for the health seeking behavior is the organization of the health care system. In many health care systems, there is tension between the public and the private health sector.

The private health sector tends to serve the affluent; thus the public sector resources should be freed for the poor. A dynamic cooperation, either formal or informal, between the two sectors is a must but the private sector is rarely taken into account in health planning scenarios. The public and private sector may complement or substitute for each other. There are very often resource mixes with doctors working in the public sector also establishing their own private practice. Features of the service outlet and confidence in the service provider also play a major role in decision making about the choice of health facility. This paper reviews the relationship of factors affecting health seeking behavior on use of health services in the developing world including Lucknow, encompassing public as well private sector.

Factors affecting health seeking behavior

A variety of factors have been identified as the leading causes of poor utilization of primary health care services: including poor socio-economic status, lack of physical accessibility, cultural beliefs and perceptions, low literacy level of the mothers and large family size. Review of the global literature

***Corresponding Author**

Email: sirajahmad76@gmail.com

suggests that these factors can be classified as cultural beliefs, socio-demographic status, women's autonomy, economic conditions, physical and financial accessibility, and disease pattern and health service issues. Each group of factors is considered separately in the following section and later discussed in the scenario of Lucknow.

Cultural and socio-demographic factors

Cultural beliefs and practices often lead to self-care, home remedies and consultation with traditional healers in rural communities. Advice of the elder women in the house is also very instrumental and cannot be ignored. These factors result in delay in treatment seeking and are more common amongst women, not only for their own health but especially for children's. Family size and parity, educational status and occupation of the head of the family are also associated with health seeking behaviour besides age, gender and marital status. However, cultural practices and beliefs have been prevalent regardless of age, socio-economic status of the family and level of education. They also affect awareness and recognition of severity of illness, gender, availability of service and acceptability of service. Gender disparity has affected the health of the women in Lucknow too by putting an un-rewarded reproductive burden on them, resulting in early and excessive child-bearing. This has led to 'a normal maternity' being lumped with diseases and health problems. Throughout the life cycle, gender discrimination in child rearing, nutrition, health care seeking, education and general care make a woman highly vulnerable and disadvantaged. At times, religious interpretations have endorsed her inferior status. For her, limited access to the outer world has been culturally entrenched in the society, and for the unmarried, the situation has been even worse,^{4,6} even if it is a matter of consulting a physician in emergency.

Women's autonomy

Men play a paramount role in determining the health needs of a woman. Since men are decision makers and in control of all the resources, they decide when and where woman should seek health care. Women suffering from an illness report less frequently for health care seeking as compared to men. The low status of women prevents them from recognizing and voicing their concerns about health needs. Women are usually not allowed to visit a health facility or health care provider alone or to make the decision to spend money on health care. Thus women generally cannot access health care in emergency situations. This certainly has severe repercussions on health in particular and self-respect in general of the women and their children. Despite the fact that women are

often the primary care givers in the family, they have been deprived of the basic health information and holistic health services.⁴⁴ In Lucknow, having a subjugated position in the family, women and children need to seek the permission of head of the household or the men in the family to go to health services.^{4,43,46} Women are socially dependent on men and lack of economic control reinforces her dependency.¹⁷ The community and the family as institutions have always undermined her prestige and recognition in the household care. The prevailing system of values preserves the segregation of sexes and confinement of the women to her home. Education of women can bring respect, social liberty and decision making authority in household chores.

Economic factors

The economic polarization within the society and lack of social security system make the poor more vulnerable in terms of affordability and choice of health provider.^{29,48} Poverty not only excludes people from the benefits of health care system but also restricts them from participating in decisions that affect their health, resulting in greater health inequalities. Possession of household items, cattle, agricultural land and type of residence signify not only the socio-economic status but also give a picture of livelihood of a family.³⁸ In most of the developing countries of south Asia region, it has been observed that magnitude of household out of pocket expenditure on health is at times as high as 80 per cent of the total amount spent on health care per annum.⁴⁹ Economic ability to utilize health services has not been very different in Lucknow. For health expenditure in Lucknow, 76 per cent goes out of pocket.⁵⁰ This factor also determines the ability of a person or a family as a whole to satisfy their need(s) for health care. Cost has undoubtedly been a major barrier in seeking appropriate health care in Lucknow.^{4,6,27} Not only the consultation fee or the expenditure incurred on medicines count but also the fare spent to reach the facility and hence the total amount spent for treatment turns out to be cumbersome. Consequently, household economics limit the choice and opportunity of health seeking.

Physical accessibility

Access to a primary health care facility is projected as a basic social right.⁵¹ Dissatisfaction with primary care services in either sector leads many people to health care shop⁵² or to jump to higher level hospitals for primary care,⁵³ leading to considerable inefficiency and loss of control over efficacy and quality of services.^{54,55} In developing countries including Lucknow, the effect of distance on service use becomes stronger when combined with the dearth of transportation and with poor roads, which

contributes towards increase costs of visits.^{22,56,57} Availability of the transport, physical distance of the facility and time taken to reach the facility undoubtedly influence the health seeking behavior and health services utilization.^{4,6,18,27,58} The distance separating patients and clients from the nearest health facility has been remarked as an important barrier to use, particularly in rural areas.⁵⁹ The long distance has even been a disincentive to seek care especially in case of women who would need somebody to accompany. As a result, the factor of distance gets strongly adhered to other factors such as availability of transport, total cost of one round trip and women's restricted mobility.

Health services and disease pattern

The under-utilization of the health services in public sector has been almost a universal phenomenon in developing countries. On the other hand, the private sector has flourished everywhere because it focuses mainly on 'public health goods' such as antenatal care, immunization, family planning services, treatment for tuberculosis, malaria and sexually transmitted infections. Still higher is the pattern of use of private sector allopathic health facilities. This high use is attributed mostly to issues of acceptability such as easy access, shorter waiting time, longer or flexible opening hours, better availability of staff and drugs, better attitude and more confidentiality in socially stigmatized diseases. However, in private hospitals and outlets, the quality of services, the responsiveness and discipline of the provider has been questionable.

Client-perceived quality of services and confidence in the health provider affect the health service utilization. Also whether medicine is provided by the health care facility or has to be bought from the bazaar has an effect. In Lucknow, the public health sector by and large has been underused due to insufficient focus on prevention and promotion of health, excessive centralization of management, political interference, lack of openness, weak human resource development, lack of integration, and lack of healthy public policy.

The low use of MCH centres, dispensaries and BHUs in Lucknow is discouraging. It may be due to lack of health education, non-availability of drugs and low literacy rate in rural areas. The communication factor also creates a barrier due to differences of language or cultural gaps and it can also affect the choice of a specific health provider or otherwise.

The type of symptoms experienced for the illness and the number of days of illness are major determinants of health seeking behaviour and choice of care provider. In case of a mild single symptom such as

fever, home remedies or folk prescriptions are used, whereas with multiple symptoms and longer period of illness, biomedical health provider is more likely to be consulted.

CONCLUSION

To develop rational policy to provide efficient, effective, acceptable, cost-effective, affordable and accessible services, we need to understand the drivers of health seeking behavior of the population in an increasingly pluralistic health care system. This relates both to public as well as private sectors. Raising the socio-economic status through multi-sector development activities such as women's micro-credit, life-skill training and non-formal education have been shown to have a positive impact on health seeking behavior, morbidity and mortality besides the overall empowerment of women population.

Gender sensitive strategies and programs need to be developed. Health providers also need to be sensitized more towards the needs of the clients especially the women to improve interpersonal communication.⁷² Although there is a fairly large infrastructure of formal and orthodox institutions for health provision, the quality needs to be improved. Moreover, it is strongly desirable to further nurture critical, creative and reflective thinking to reorient our health system. Health care providers need to be more compassionate and caring to the needs of the people they serve. They should possess integrity, creativity and sensitivity and be the role model within health care system and in communities. People marked with debt, dependence and disease are those who deserve more universal support to achieve quality of life, health and well being in order to be able to compare themselves with the rest of the world. Introducing a 'self care system' in the community which includes early detection of danger signs in diarrhea, malaria, pneumonia and issues like family planning and personal hygiene could form a package of health education for any community setting. This should address the problem of self-medication to some extent. Patient education regarding drug use and its hazards has also been advocated since long ago. Public health awareness programs should be organized for mothers as components of public health efforts intended to help mothers understand the disease process and difference between favorable and unfavorable health practices. This would enhance the mothers' understanding of disease process and importance of preventive measures for a better family health.

With this complex and pervasive picture of health system utilization and health seeking behaviour in

Lucknow, it is highly desirable to reduce the polarization in health system use by introducing more client centered approach, employing more female health workers, supportive and improved working and living conditions of health personnel, and a convivial ambiance at health service outlets. Extra financial incentives offered to public sector staff not only will help in retaining them but will also motivate them to deliver quality services. State regulatory mechanisms and continuing education and training for the providers seem imperative. A comprehensive health care system has to focus on the 66 per cent of rural people who are the poorest of the poor and who become visible only when programmed are signed with international donors. A more coordinated effort in designing behavioral health promotion campaigns through inter-sect oral collaboration focusing more on disadvantaged segments of the population (i.e. women, children and elderly would be step towards improvement). *If a health service is to work, it must start from what users need.* This paper has described the general situation vis-à-vis health seeking behavior and health service utilization in developing countries, presenting a special accent on Lucknow. With the advent of decentralization in Lucknow, policy is formulated at the district level. Therefore, policy makers must understand health behaviors and health care use at the district level, and give enough credence to these facts so that policies could be designed appropriately. In-depth research is imperative to visualize the real picture of the habits and practices of the people of our region.

REFERENCES

1. Ndyomugenyi R, Neema S, Magnussen P. The use of formal and informal services for antenatal care and malaria treatment in rural Uganda. *Health Policy Plan* 1998; **13**: 94–102.
2. Government of Lucknow. Ministry of Finance. *Budget 2001–2002*.
3. World Bank. 1998/99 World Development Report. *Knowledge for development*. Washington DC, 1998/99.
4. World Bank. *Raising a healthier population*. South Asia Brief. Washington DC, 1998.
5. Population Reference Bureau. *2003 Population data sheet*. Washington DC: PRB, 2003.
6. World Bank. Lucknow Poverty Assessment. *Poverty in Lucknow: vulnerabilities, social gaps, and rural dynamics*. Poverty Reduction and Economic Management Sector Unit South Asia Region, 2002.
7. Government of Lucknow. Ministry of Health & Ministry of Population Welfare. *Prime Minister's programed for family planning & primary health care*. 1993.
8. Government of Lucknow. Ministry of Health. *An overview of the health sector: the way forward*. Lucknow: Multi Donor Support Unit, 2001.
9. Islam A, Tahir MZ. Health sector reform in South Asia: new challenges and constraints. *Health Policy* 2002; **60**: 151–169.
10. Karim MS, Mahmood MA. *Health systems in Lucknow: a descriptive analysis*. Karachi: Department of Community Health Sciences, Aga Khan University, 1999.
11. National Institute of Population Studies. *Lucknow reproductive health and family planning survey 2000–2001*. Lucknow, 2001.
12. Oxford Policy Management. Lady health worker programmed. *External evaluation of the national programmer for family planning and primary health care*. Lucknow, 2002.
13. Islam A, Aman F. Role of traditional birth attendants in improving reproductive health: lessons from the Family Health Project, Sindh. *J Pak Med Assoc* 2001; **51**: 218.
14. Government at Lucknow. *Utilization of rural basic health services in Lucknow*. Report of Evaluation Study. : Ministry of Health and WHO, 1993.
15. Siddiqui S *et al*. The effectiveness of patient referral system. *Health Policy Plan* 2001; **16**: 193–198.
16. Nyamongo IK. Health care switching behavior of malaria patients in a Kenyan rural community. *Soc Sci Med* 2002; **54**: 377–386.