

**INTERNATIONAL JOURNAL OF ENGINEERING SCIENCES & MANAGEMENT****AN APPLICATION OF FRACTIONAL CALCULUS IN SOLVING NEW MODIFIED GENERALIZED FRACTIONAL KINETIC EQUATION****Manoj Sharma**

\* Department of Mathematics RJIT, BSF Academy, Tekanpur

**ABSTRACT**

In this paper, we derive the solution of new generalized fractional kinetic equation involving the New generalized Miller-Ross function. The result obtained here is quite general in nature and capable of yielding a very large number of results. Special cases, involving the M-Series, Mittag-Leffler function and Miller-Ross function etc. are also considered.

Mathematics Subject Classification: 33C60, 82C31

**Keywords:** Fractional kinetic equation, new generalized Miller-Ross function, Mittag-Leffler function, Riemann-Liouville operator, Laplace transform, generalized M-series.

**INTRODUCTION**

The great importance of mathematical physics in distinguished astrophysical problems has attracted astronomers, mathematician and physicists to pay more attention to available mathematical tools that can be widely used in solving several problems of physics, applied physics and astrophysics. A spherically symmetric non-rotating, self-gravitating model of star like the Sun is assumed to be in thermal equilibrium and hydrostatic equilibrium. The star is characterized by its mass, luminosity, effective surface temperature, radius, central density and central temperature. The stellar structures and their mathematical models are investigated on the basis of above characters and some additional information related to the equation of state, nuclear energy generation rate and the opacity. The assumptions of thermal equilibrium and hydrostatic equilibrium imply that there is no time dependence in the equations describing the internal structure of the star (Kourganoff [11], Perdang [17] and Clayton [2]). Energy in such stellar structures is being produced by the process of chemical reactions (thermonuclear reactions).

Consider an arbitrary reaction characterized by a time dependent quantity  $N = N(t)$ . It is possible to calculate rate of change  $dN/dt$  to a balance between the destruction rate  $d$  and the production rate  $p$  of  $N$ , that is  $\frac{dN}{dt} = -d + p$ . In general, through feedback or other interaction mechanism, destruction and production depend on the quantity  $N$  itself:  $d = d(N)$  or  $p = p(N)$ . This dependence is complicated since the destruction or production at time  $t$  depends not only on  $N(t)$  but also on the past history  $N(\tau)$ ,  $\tau < t$ , of the variable  $N$ .

This may be formally represented by (Haubold and Mathai [7])

$$\frac{dN}{dt} = -d(Nt) + p(Nt), \quad (1)$$

where  $Nt$  denotes the function defined by  $Nt(t*) = N(t - t*)$ ,  $t* > 0$ .

Haubold and Mathai [7] studied a special case of this equation, when spatial fluctuations or inhomogenities in quantity  $N(t)$  are neglected, is given by the equation

$$\frac{dN_i}{dt} = -c_i N_i(t) \quad (2)$$

with the initial condition that  $N_i(t = 0) = N_0$  is the number density of species  $i$  at time  $t = 0$ ; constant  $c_i > 0$ , known as standard kinetic equation.

The solution of the equation (2) is given by

$$N_i(t) = N_0 e^{-c_i t} \quad (3)$$

An alternative form of the same equation can be obtained on integration:

$$N(t) - N_0 = c_0 D_t^{-1} N(t) \quad (4)$$

where  ${}_0D_t^{-1}$  is the standard integral operator. Haubold and Mathai [7] have given the fractional generalization of the standard kinetic equation (2) as

$$N(t) - N_0 = c^v {}_0D_t^{-v} N(t) \quad (5)$$

where  ${}_0D_t^{-\nu}$  is the well known Riemann-Liouville fractional integral operator (Oldham and Spanier [16]; Samko et al. [19]; Miller and Ross [13]) defined by

$${}_0D_t^{-\nu}N(t) = \frac{1}{\Gamma(\nu)} \int_0^t (t - u)^{\nu-1} f(u) du, \quad R(\nu) > 0, \tag{6}$$

The solution of the fractional kinetic equation (6) is given by (see Haubold and Mathai [7])

$$N(t) = N_0 \sum_{k=0}^{\infty} \frac{(-1)^{vk}}{\Gamma(\nu k + 1)} (ct)^{\nu k} \tag{7}$$

Further Saxena, Mathai and Haubold [20] studied the generalizations of the fractional kinetic equation in terms of the Mittag-Leffler functions which extended the work of Haubold and Mathai [7]. In an another paper Saxena, Mathai and Haubold [21] developed the solutions for fractional kinetic equations associated with the generalized Mittag-Leffler function and R-function.

In the present paper we introduce and investigate the further computable extensions of the generalized fractional kinetic equation. The fractional kinetic equation and its solution, discussed in terms of the generalized Miller-Ross, are written in compact and easily computable form.

**THE NEW MODIFIED GENERALIZED MILLER-ROSS FUNCTION**

This function introduced by the authors is defined as follows:

$$N_{p,q}^{\alpha,\beta,a} (a_1 \dots a_p, \gamma, \delta; , b_1 \dots b_q, \rho; x ) = \sum_{n=0}^{\infty} \frac{(a_1)_{nk} \dots (a_p)_{nk} (\gamma)_{nk} (\delta)_{nk} a^n x^{n+\beta}}{(b_1)_{nk} \dots (b_q)_{nk} (\rho)_{nk} \Gamma(\alpha n + \beta + 1)} \tag{8}$$

Here,  $p$  upper parameters  $a_1, a_2, \dots, a_p$  and  $q$  lower parameters  $b_1, b_2, \dots, b_q, \alpha, \beta \in C, R(\alpha) > 0, R(\beta) > 0$  and  $(a_j)_{nk}, (b_j)_{nk}, (\gamma)_{nk}, (\delta)_{nk}, (\rho)_{nk}$  are pochhammer symbols. The function (3) is defined when none of the denominator parameters  $b_j$ s,  $j = 1, 2, \dots, q$  is a negative integer or zero. If any parameter  $a_j$  is negative then the function (3) terminates into a polynomial in  $x$ . By using ratio test, it is evident that function (3) is convergent for all  $x$ , when  $q \geq p$ , it is convergent for  $|x| < 1$  when  $p = q + 1$ , divergent when  $p > q + 1$ . In some cases the series is convergent for  $x = 1, x = -1$ . Let us consider take,

$$\beta = \sum_{j=1}^p a_j - \sum_{j=1}^q b_j$$

when  $p = q + 1$ , the series is absolutely convergent for  $|x| = 1$  if  $R(\beta) < 0$ , convergent for  $x = -1$ , if  $0 \leq R(\beta) < 1$  and divergent for  $|x| = 1$ , if  $1 \leq R(\beta)$ .

**GENERALIZED FRACTIONAL KINETIC EQUATIONS**

'In this section we investigate the solution of generalized fractional kinetic equation'. The results are obtained in a compact form in terms of New Generalized Miller-Ross function and are suitable for computation. The result is presented in the form of a theorem as follows:

**Theorem 1:**

If  $\nu > 0, c > 0, \mu > 0$ , then for the solution of the generalized fractional kinetic equation

$$N(t) - N_0 t^{-\mu(\nu-1)} N_{p,q}^{\nu,\mu,a} (a_1 \dots a_p, \gamma, \delta; , b_1 \dots b_q, \rho; t^\nu ) = -c^\nu {}_0D_t^{-\nu} N(t) \tag{9}$$

there holds the formula

$$N(t) = N_0 t^{-\mu(\nu-1)} \sum_{k=0}^{\infty} (-1)^k (ct)^{k\nu} N_{p,q}^{2\nu,\mu,a} (a_1 \dots a_p, \gamma, \delta; , b_1 \dots b_q, \rho; t^\nu ) \tag{10}$$

**Proof.** Applying the Laplace transform both the sides of equation (9), we get

$$L\{N(t)\} - L\{N_0 t^{-\mu(v-1)} N_{p,q}^{v,\mu,a}(a_1 \dots a_p, \gamma, \delta; b_1 \dots b_q, \rho; t^v)\} = L\{-c^v {}_0D_t^{-v} N(t)\}$$

$$N(s) - N_0 \sum_{k=0}^{\infty} \frac{(a_1)_{nk} \dots (a_p)_{nk} (\gamma)_{nk} (\delta)_{nk} a^k}{(b_1)_{nk} \dots (b_q)_{nk} (\rho)_{nk} s^{vk+\mu+1}} = -c^v s^{-v} N(s) \tag{11}$$

Solving for  $N(s)$ , it gives

$$N(s) = \frac{N_0}{(1 + c^v s^{-v})} \sum_{k=0}^{\infty} \frac{(a_1)_{nk} \dots (a_p)_{nk} (\gamma)_{nk} (\delta)_{nk} a^k}{(b_1)_{nk} \dots (b_q)_{nk} (\rho)_{nk} s^{vk+\mu+1}} \tag{12}$$

Now, taking inverse Laplace transform both the sides of (12), we get

$$L^{-1}\{\bar{N}(s)\} = L^{-1}\left\{N_0 \sum_{k=0}^{\infty} \frac{(-1)^k (c^v s^{-v})^k (1)_k}{k!} \sum_{k=0}^{\infty} \frac{(a_1)_{nk} \dots (a_p)_{nk} (\gamma)_{nk} (\delta)_{nk} a^k}{(b_1)_{nk} \dots (b_q)_{nk} (\rho)_{nk} s^{vk+\mu+1}}\right\} \tag{13}$$

Or

$$N(t) = N_0 t^{-\mu(v-1)} \sum_{k=0}^{\infty} (-1)^k (ct)^{kv} N_{p,q}^{2v,\mu,a}(a_1 \dots a_p, \gamma, \delta; b_1 \dots b_q, \rho; t^v) \tag{14}$$

This is complete proof of the statement (9).

**SPECIAL CASES**

When  $a = 1, \mu = 0$ , then

**Corollary: 1.** If  $v > 0, c > 0, \mu > 0$ , then for the solution of the generalized fractional kinetic equation (in form of M-Series [22, 23])

$$N(t) - N_0 M_{p,q}^v(a_1 \dots a_p, \gamma, \delta; b_1 \dots b_q, \rho; t^v) = -c^v {}_0D_t^{-v} N(t) \tag{15}$$

there holds the formula

$$N(t) = N_0 \sum_{k=0}^{\infty} (-1)^k (ct)^{kv} M_{p,q}^{2v}(a_1 \dots a_p, \gamma, \delta; b_1 \dots b_q, \rho; t^v) \tag{16}$$

When  $\mu = 0, a = 1$  and no upper and lower parameter then

**Corollary: 2.** If  $v > 0, c > 0, \mu > 0$ , then for the solution of the generalized fractional kinetic equation (in form of Mittag-Leffler function [10])

$$N(t) - N_0 E_v(t^v) = -c^v {}_0D_t^{-v} N(t) \tag{17}$$

there the relation

$$N(t) = N_0 \sum_{k=0}^{\infty} (-1)^k (ct)^{kv} E_v(t^v) \tag{18}$$

When  $v = 1$  and no upper and lower parameter then

**Corollary: 3.** If  $v > 0, c > 0, \mu > 0$ , then for the solution of the generalized fractional kinetic equation (in form of Miller-Ross function [27])

$$N(t) - N_0 E_t(\mu, a) = -c^1 {}_0D_t^{-1} N(t) \tag{19}$$

there exist the result

$$N(t) = N_0 \sum_{k=0}^{\infty} (-1)^k (ct)^k E_t(\mu, a) \tag{20}$$

**CONCLUSION**

In this paper we have introduced an extended fractional generalization of the standard kinetic equation and established solution for the same. Fractional kinetic equation can be used to compute the particle reaction rate and describes the statistical mechanics associated with the particle distribution function. The generalized fractional kinetic equation discussed in this article, involving New Generalized Miller-Ross function contains a number of known (may be new also) fractional kinetic equations involving various other special functions (the M-series, Mittag-Leffler function etc.).

**REFERENCE**

1. R.P. Agarwal, A propos d'une note de M.Pierre Humbert, C.R. Acad. Sci. Paris 236 (1953), 2031-2032.
2. D.D. Clayton, Principles of Stellar Evolution and Nucleosynthesis, 2nd ed.: Chicago: Chicago Univ. Chicago Press, 1983.
3. A. Erdélyi, W. Magnus, F. Oberhettinger and F.G. Tricomi, Higher Transcendental Functions, Vol.1, McGraw-Hill, New York-Toronto-London, 1953.
4. A. Erdélyi, W. Magnus, F. Oberhettinger and F.G. Tricomi, Tables of Integral Transform, Vol.1, McGraw-Hill, New York-Toronto-London, 1954.
5. A. Erdélyi, W. Magnus, F. Oberhettinger and F.G. Tricomi, Higher Transcendental Functions, Vol.3, McGraw-Hill, New York-Toronto-London, 1955.
6. F. Ferro, A. Lavago and P. Quarati, Temperature dependence of modified CNO nuclear reaction rates in dense stellar plasmas, arXiv:nuc1h/ 0312106v1, 2003.
7. H.J. Haubold and A.M. Mathai, The fractional kinetic equation and thermonuclear functions, Astrophysics and Space-Sciences 327 (2000), 53-63.
8. R. Hilfer (ed.), Applications of Fractional Calculus in Physics, World Scientific, Singapore, 2000.
9. P. Humbert, Quelques resultats relatifs a la fonction de Mittag-Leffler, C.R. Acad. Sci., Paris 236 (1953), 1467-1468.
10. P. Humbert and R.P. Agarwal, Sur la fonction de Mittag-Leffler et quelques-unes de ses generalizations, Bull. Sci. Math. (Ser. II) 77 (1953), 180-185.
11. V. Kiryakova., some special functions related to fractional calculus and fractional (non-integer) order control systems and equations. Facta Universitatis (Sci. J. of Univ. Nis) Automatic Control and Robotics, 7 No.1 (2008), 79-98.
12. K.R. Lang, Astrophysical Formulae Vol.1 (Radiation, Gas Processes and High Energy Astrophysics) and Vol.II (Space, Time, Matter and Cosmology), Springer-Verlag, Berlin-Heidelberg, 1999.
13. K.S. Miller and B. Ross, An Introduction to the Fractional Calculus and Fractional Differential Equations, John Wiley and Sons, New York, 1993.
14. G.M. Mittag-Leffler, Sur la nouvelle fonction  $E_\alpha(x)$ , C.R. Acad. Sci., Paris (Ser.II) 137 (1903), 554-558.
15. G.M. Mittag-Leffler, Sur la representation analytique d'une branche uniforme d'une fonction monogene, Acta Math. 29 (1905), 101-181.
16. K.B. Oldham and J. Spanier, The Fractional Calculus: Theory and Applications of Differentiation and Integration to Arbitrary Order, Academic Press, New York, 1974.
17. J. Perdang, Lecture Notes in Stellar Stability, Part I and II, Instituto di Astronomia, Padova, 1976.
18. T.R. Prabhakar, A singular integral equation with generalized Mittag-Leffler function in the kernel, Yokohama Mathematical Journal 19 (1971), 7-15.
19. S.G. Samko, A.A. Kilbas and O.I. Marichev, Fractional Integrals and Derivatives: Theory and Applications, New York, 1993.
20. R.K. Saxena, A.M. Mathai and H.J. Haubold, On fractional kinetic equations, Astrophysics Space Sci. 282 (2002), 281-287.
21. R.K. Saxena, A.M. Mathai and H.J. Haubold, On generalized fractional kinetic equations, Physica A 344 (2004), 653-664.
22. M. Sharma and R. Jain, A note on a generalized M-series as a special function of fractional calculus, Fract. Calc. Appl. Anal. 12 No.4 (2009) 449-452.
23. M. Sharma, Fractional integration and fractional differentiation of the M-series, Fract. Calc. Appl. Anal. 11 No.2 (2008), 187-192.
24. H.M. Srivastava and R.K. Saxena, Operators of fractional integration and their applications, Applied Mathematics and Computation 118 (2001), 1- 52.

25. C. Tsallis, Entropic nonextensivity: A possible measures of Complexity, Chaos, Solitons and Fractals 13 (2002), 371-391.
26. A. Wiman, Ueber den Fundamentalsatz in der Theorie der Funkilionen  $E_\alpha(x)$ , Acta. Math. 29 (1905), 191-201.
27. Carl. F. Lorenzo, Tom T. Hartley, Generalized Functions for the Functional Calculus, Nasa/tp-1999-209424/rev 1, p1-17.